

Promoters

Prof. dr. Pierre Dorny
Faculty of Veterinary Medicine, Ghent University;
Institute of Tropical Medicine, Antwerp

Prof. dr. Dirk Berkvens
Faculty of Bioscience Engineering, Ghent
University
Institute of Tropical Medicine, Antwerp

Dr. Dirk Geysen
Institute of Tropical Medicine, Antwerp

Prof. dr. Gobena Ameni
Aklilu Lemma Institute of Parasitology, Addis
Ababa University, Ethiopia

Members of the Examination Committee

Prof. dr. Luc Duchateau
Faculty of Veterinary Medicine, Ghent University
Chairman of the Examination Committee

Prof. dr. Bruno Levecké
Faculty of Veterinary Medicine, Ghent University
Secretary of the Examination Committee

Prof. dr. Edwin Claerebout
Faculty of Veterinary Medicine, Ghent University,

Prof. dr. Bertrand Losson
Faculty of Veterinary Medicine, University of
Liège

Dr. Nick De Regge
Sciensano

Curriculum Vitae

Teshale Sori Tolera holds a DVM (1999) from Addis Ababa University and a MSc degree (2010) from the Institute of Tropical Medicine, Belgium and has attended several face-to-face and online postgraduate courses in veterinary epidemiology and biostatistics.

After working for two years (2000-2001) as an assistant researcher at Bako Agricultural Research Centre (Ethiopia) and 2.5 years (2002-2004) as a junior research officer at the National Veterinary Institute at Bishoftu (Ethiopia), he joined the College

of Veterinary Medicine and Agriculture, Addis Ababa University in 2004 where he was appointed as a lecturer. He teaches analytical veterinary epidemiology, large animal medicine and infectious diseases of poultry and has been involved in several research activities and MSc and DVM coaching. His research focuses on the impact of ticks and tick-borne diseases on the national economy and public health, the dynamics of tick-borne diseases in various epidemiological situations. Recently he has been involved in the design, implementation and monitoring of intervention studies of Newcastle Disease and Infectious Bursal Disease and was asked to strengthen the interaction of research and clinical aspects of Newcastle disease. He has produced several publications in peer reviewed international journals.



INVITATION

Public Defence of the doctoral thesis of

Teshale Sori Tolera

May 16th 2018

Laboratory of Parasitology
Department of Virology, Parasitology and
Immunology
Faculty of Veterinary Medicine, UGent



FACULTEIT DIERGENEESKUNDE
approved by EAEVE

You are kindly invited to attend the public defence
of the doctoral thesis of

Teshale Sori Tolera

Title of the thesis:

**Study on Tick-Borne Pathogens
and Tick-Borne Diseases Using
Molecular Tools with Emphasis
on *Anaplasma* spp. and *Ehrlichia*
spp. In Ticks and Domestic
Ruminants in Ethiopia**

The public defence will take place on

Wednesday 16th May 2018

at 17:00 hours

in *kliniekauditorium Hoogbouw* - Entrance 24 of

the Faculty of Veterinary Medicine, UGent

Salisburylaan 133, Merelbeke

**You are cordially invited to a reception that
will be held after the public defence**

Summary of the Thesis

Although tick-borne pathogens are among the most important constraints in the domestic ruminant industry causing huge economic losses, little information is available on their occurrence and distribution in Ethiopia. Within this doctoral study, three cross-sectional epidemiological studies using molecular techniques were conducted, one on ticks collected from cattle and sheep, one on domestic ruminants and one on unfed ticks collected from the field.

The survey conducted on ticks collected from cattle and sheep using a molecular analysis of 18S rDNA identified the occurrence of *Theileria buffeli/orientalis*, *Theileria velifera*, and *Theileria ovis* in *Rhipicephalus evertsi evertsi* and *Rhipicephalus decoloratus*. According to the 16s rDNA PCR and sequencing, six species of *Bartonella*, three species of *Rickettsia* (*Rickettsia africae*, *Rickettsia felis* and *Rickettsia* sp.), *Anaplasma ovis*, *Ehrlichia ruminantium*, *Ehrlichia* spp., *Anaplasma* spp. and *Borrelia burgdorferi* s.l. were identified in *Rhipicephalus* spp. This is the first report of the occurrence of *A. ovis*, *Bartonella elizabethae*, *Bartonella bovis*, *Bartonella koehlerae*, *Bartonella quintana* and *Bartonella vinsonii berkhoffii* in Ethiopia. Our finding highlights the risk of infection of animals and humans with zoonotic tick-borne bacteria in Ethiopia. At the end of the first survey, we focused on the molecular identification of *Ehrlichia* spp. and *Anaplasma* spp., in which we developed a semi-nested PCR amplifying 925bp of 16S rDNA. A pair of primers designated EBR2 and EBR3 was designed from the *Anaplasma* 16S rDNA sequences for simultaneous detection of *Ehrlichia* spp. and *Anaplasma* spp. Individual species of *Ehrlichia* and *Anaplasma* were identified by restriction with *MbolI*, *HhaI* and *MspI* enzymes, including mixed infections. Analysis of *Amblyomma* spp. from various parts of Ethiopia for bacterial pathogens using this molecular method revealed the occurrence of *Anaplasma marginale*, *Anaplasma phagocytophilum*, *Anaplasma centrale*, *Anaplasma* (formerly *Ehrlichia*) sp. *Omatjenne* and *E. ruminantium*. By helping animal health professionals in their decision-making regarding the diagnosis and control of anaplasmosis and heartwater, this improved molecular method will help the government in establishing a disease-monitoring program in the dairy and beef industries.

The second part involves surveys of *A. phagocytophilum* and *A. sp. Omatjenne* infection in cattle in Africa and that of *Anaplasma* spp. and *Ehrlichia* spp. in cattle, sheep and goats in Ethiopia. In the international survey, respectively 19 (2.7%) and 45 (6.5%) samples yielded positive signals for *A. phagocytophilum* and *A. sp. Omatjenne*. *Anaplasma* sp. *Omatjenne* was detected in all countries except Tanzania while *A. phagocytophilum* was detected only in samples originating from Ethiopia. Out of the 922 blood samples from cattle, sheep and goats from five different localities in Ethiopia analyzed by 16S rDNA PCR, 523 (56.7%) tested positive. Overall, 67.4% of cattle, 65% of sheep and 18%

of goats tested positive for one or more *Anaplasma* spp. No positive result was observed for *Ehrlichia* spp. using the 16S rDNA analysis. RFLP analysis identified *A. marginale*, *A. ovis*, *A. phagocytophilum*, *A. centrale* and *A. sp. Omatjenne*. We conclude that infection of domestic ruminants with *Anaplasma* spp. is widespread in Ethiopia. Livestock improvement plans through introduction of improved breeds should be aware of this and take the necessary precautions to minimize losses associated with anaplasmosis. First reports of infection of domestic ruminants with *A. phagocytophilum*, *A. ovis* and *A. sp. Omatjenne* are provided for Ethiopia. In addition, the occurrence of *A. phagocytophilum* together with *A. sp. Omatjenne* outside Europe and South Africa is made for the first time. Random samples of 493 animals were tested with a pCS20 PCR for the identification of *E. ruminantium*. Three samples from 75 (4%) cattle gave a positive result. *Ehrlichia ruminantium* was identified in five of 13 clinical cases of heartwater in dairy cows (38.46%) and 30 of 72 (41.67%) clinically affected Boer goats.

Identification of ticks that carry pathogens of veterinary and public-health importance, belonging to the genus *Anaplasma*, is an initial step to identify tick vectors that play a role in the epidemiology of the diseases they cause. A molecular investigation was carried out to identify *Anaplasma* spp. carried by unfed field ticks with an emphasis on *A. phagocytophilum* and *A. ovis*. A total of 240 unfed ticks (adults and nymphs) were collected and analyzed using PCR-RFLP. *Anaplasma ovis* was identified in *R. evertsi*, *Amblyomma* spp. and *Hyalomma* spp. *Anaplasma phagocytophilum* was detected only in *Rhipicephalus pulchellus*. Our finding identified potential vectors of *A. ovis* to be further confirmed by experimental study.

In conclusion, this doctoral study identified tick-borne pathogens of veterinary and public health importance in Ethiopia. An improved molecular method was developed for simultaneous detection of *Anaplasma* spp. and *Ehrlichia* spp. and it was used in epidemiological studies in domestic ruminants and ticks. A high prevalence of anaplasmosis and a low prevalence of heartwater were observed. The occurrence of infection in cattle with *A. sp. Omatjenne* was identified in certain African countries. Furthermore, candidate vectors of *A. ovis* were identified. Further studies on the medical significance of the identified zoonotic pathogens, search for vectors of *A. phagocytophilum* and experimental studies for the confirmation of the vectors of *A. ovis* are warranted. Lastly, an improvement of molecular methods is needed for the detection of carriers of *E. ruminantium*.